

Original Article

Production of Tomato Seedlings using Seeds Pelleted with *Natrum muriaticum* and Submitted to Saline Stress

Filipe Pereira Giardini Bonfim¹, Vicente Wagner Dias Casali², Aline Mako Yoshikawa^{1*}

1 – Universidade Júlio de Mesquita Filho - UNESP, SP, Brasil.

2 – Universidade Federal de Viçosa, MG, Brasil.

*alinemako8@hotmail.com – <https://orcid.org/0000-0002-3197-9253>

Abstract

The development and yield of plants are directly related to the effects of salinity. Several scientific studies report a significant reduction in the growth and production of tomatoes in soils with high electrical conductivity. The correction or recovery of salinized soils, although technically possible, is a slow and costly process, making it necessary to develop new technologies. This study aims to evaluate the production of tomato (*Solanum lycopersicon* L.) seedlings using seeds pelleted with a homeopathic preparation of *Natrum muriaticum* (*Nat-m*) submitted to saline stress. The treatments consisted of the pelletization of tomato seeds with six dynamizations of *Nat-m* (3cH, 5cH, 7cH, 9cH, 11cH, and 13cH). Coated or uncoated seeds (controls) were placed in phenolic foam blocks, kept in plastic trays previously moistened with 50 mM NaCl saline solution (2.922 g.L⁻¹ of NaCl, electrical conductivity = 4.5 dSm⁻¹) and nutrient solution (0.15 dS.m⁻¹) at half the ionic strength, containing 4, 1, 2, 1, 0.5, and 0.5 mmol.L⁻¹ of N, P, K, Ca, Mg and S, and 17.5, 9.5, 10.5, 2, 0.45 and 0.35 mmol.L⁻¹ of Fe, Mn, Zn, Cu, and Mo, respectively. The variables evaluated were the percentage of emerged plants, emergence speed index, shoot length, leaf area, number of leaves, root volume, root dry matter, and shoot dry matter. The treatments pelleted seeds/talc *Nat-m* 5cH and pelleted seeds/talc *Nat-m* 7cH increased all variables evaluated. However, they differed statistically from the controls, positively developing tomato seedlings under disequilibrium conditions.

Keywords: Salinity, homeopathic preparations, *Solanum lycopersicon*, pelletization, homeostasis.

Introduction

About 1.5 million hectares of arable land are lost each year due to the accumulation of salts. Salinization has been identified as the primary process of soil degradation. Although the information on saline areas is not well established in Brazil, it is estimated that 20-25% of irrigated areas present salinization problems [1]. In general, the soil is considered saline when the amount of salts in the soil can hinder the development of plants. In most cultures, this happens when the electrical conductivity of the saturation extract is equal to or greater than 2 dS.m⁻¹ [2].

The interaction of salts in plants occurs directly in the enzymatic activity. Consequently, physiological and biochemical processes, such as respiration, photosynthesis, and metabolism of proteins and nucleic acids, are affected [3], promoting a high energy expenditure to maintain these processes.

Plants considered sensitive to salinity do not develop or are economically viable when the electrical conductivity of the saline concentration of the medium is higher than 1.30 dS.m⁻¹. Moderately sensitive plants grow in solutions with electrical conductivity between 1.30 and 3.0 dS.m⁻¹. Plants are relatively tolerant of conductivity between 3.00 and 6.00 dS.m⁻¹, and plants tolerate a conductivity between 6.00 and 10.00 dS.m⁻¹ [4]. Tomato (*Solanum lycopersicon* L.), a vegetable extensively consumed in Brazil, is moderately sensitive to soil salinity, mainly during early development [5].

The accumulation of salts in cultivated soils is quite common, mainly due to high doses of fertilizers, lack of leaching of accumulated salts after cultivation, and poor quality irrigation water [6]. Moreover, the recovery of these soils requires high investments and demands a long time [7]. Therefore, the substitution of conventional practices for ecologically correct and economically feasible methods is necessary not to impair the environment, producers, and the final product.

The use of homeopathy as an input is an approved practice. It has been growing in consequence of advances in non-conventional food production systems. It is used in several segments of agriculture, including germination [8], seedling production [9], pest control [10], plant diseases [11], increase in active substances [12], detoxification of plants due to contamination by metals such as aluminum and copper [13], and plant metabolism [14].

Ultra-diluted substances act constructively and defensively in the vitality systems of living beings. Through self-regulation, plants respond with great intensity to homeopathy when applied in a stress situation [15]. Thus, the objective of this study was to evaluate the production of tomato seedlings (*Solanum lycopersicon* L.) using seeds pelleted with dynamizations of homeopathic preparation *Natrum muriaticum*, submitted to saline stress.

Material and Methods

Choice of treatments

Natrum muriaticum (Nat-m) 3cH, 5cH, 7cH, 9cH, 11cH, and *Natrum muriaticum* 13cH were chosen according to indications of crops that grow in soils with a tendency to salinity [15].

Preparation of inputs

Natrum muriaticum 3cH was purchased from a homeopathy pharmacy in the city of Viçosa, MG, Brazil. According to the Brazilian Homeopathic Pharmacopeia, other dynamizations were prepared based on this first dynamization at the Homeopathy Laboratory of the Department of Plant Science of the Federal University of Viçosa (DFT/UFV) [16]. Respecting the ratio of 1:100, a drop of a homeopathic substance was used in 99 drops of the medium. The medium was distilled water. It was chosen to remove any interferences from ethanol with the results. The suctioning process was performed using a DENISE 10-50 "mechanical arm" dynamize (AUTIC, São Paulo, Brazil). *Natrum muriaticum* (Nat-m) 3cH, 5cH, 7cH, 9cH, 11cH and *Natrum muriaticum* 13cH were prepared before the coating of seeds.

Seed pelleting

The seed coating process, pelletization, followed an impregnation methodology respecting the weight/volume ratio (1:1). The talc was the inert pharmaceutical talc (LAB Synth, Diadema, Brazil). The composition of the talc was $Mg_3Si_4O_{10}(OH)_2$, and the adhesive was the treatments (*Nat-m* 3cH, *Nat-m* 5cH, *Nat-m* 7cH, *Nat-m* 9cH, *Nat-m* 11cH, and *Nat-m* 13cH) and distilled water. The homeopathic preparations and the distilled water were impregnated in pharmaceutical talc using a petri dish to form an integral, white paste, which later coated the seeds.

Commercial tomato seeds (*Solanum lycopersicon* L.) were used without defensives (raw seeds). The cultivar was SANTA CLARA I 5300. The germination percentage was 90%, and the purity was 99.9% (lot 28111).

Conducting the experimente

The experiment was conducted in a greenhouse of the Department of Plant Sciences of the Federal University of Viçosa in the municipality of Viçosa, Zona da Mata Mineira. The geographical coordinates were 42°52' W and 42°50' W, and 20°44' S and 20°47' S, Brazil. The experimental design

was completely randomized with eight treatments and four replications. Each experimental unit consisted of ten seeds/seedlings. The treatments are:

- 1- Pelleted seeds/talc + *Nat-m* 3cH
- 2- Pelleted seeds/talc + *Nat-m* 5cH
- 3- Pelleted seeds/talc + *Nat-m* 7cH
- 4- Pelleted seeds/talc + *Nat-m* 9cH
- 5- Pelleted seeds/talc + *Nat-m* 11cH
- 6- Pelleted seeds/talc + *Nat-m* 13cH
- 7- Pelleted seeds/talc + distilled water (control 1)
- 8- Non-pelleted seeds (control 2)

Coated or uncoated seeds (controls) were placed in phenolic foam blocks (5.0x5.0x3.8 cm), kept in plastic trays (0.80x0.40x0.10 m) previously moistened with 50 mM of NaCl saline solution (2.922 g.L⁻¹ of NaCl, electrical conductivity = 4.5 dSm⁻¹) and nutrient solution (0.15 dS.m⁻¹) [17] at half the ionic strength, containing 4, 1, 2, 1, 0.5 and 0.5 mmol.L⁻¹ of N, P, K, Ca, Mg and S, and 17.5, 9.5, 10.5, 2, 0.45, and 0.35 mmol.L⁻¹ of Fe, Mn, Zn, Cu and Mo, respectively. After 28 days of planting, the evaluations resumed.

Variables

The evaluated variables were germination percentage, germination speed index, shoot length (mm), leaf area (cm²), number of leaves, root volume (mL), root dry matter (g), and shoot dry matter (g).

The number of total normal seedlings per plot determined the germination percentage on the fourteenth day after sowing. The determination of the germination speed index was performed according to Maguire [18]. It is given by the sum of the ratio between the number of seeds germinated each day and the day of the evaluation according to the formula: $GSI = G1/N1 + G2/N2 + G3/N3 + \dots + Gn/Nn$, where: G1, G2, G3, ..., Gn is the number of seeds germinated on the day of observation and N1, N2, 3, ..., Nn is the number of days after sowing.

The shoot length was determined using a digital caliper. The leaf area was determined using a leaf area integrator (ΔT Mater MK2 Area). The root volume was determined by the difference in volume (using a 10mL beaker). Shoot dry matter of dry roots was obtained in an oven with forced ventilation at 45°C until constant weight. Data were subjected to analysis of variance, and the means were compared by Tukey test at 5% probability using the SAEG 9.0 software.

Results and Discussions

The variables evaluated were germination percentage (GP), germination speed index (GSI), shoot length (SL), leaf area (LA), number of leaves (NL), root volume (RV), root dry matter (RDM), and dry shoot matter (SDM). F test presented significant differences at 5% probability in the function of treatments, thus evidencing the action of the homeopathic preparation *Natrum muriaticum* under the experimental conditions (Tables 1 and 2).

The treatment pelleted seeds/talc + distilled water (control 1) did not differ from the treatment without pelleting (control 2) for all variables evaluated (Tables 3 and 4). Thus, there is an innocuousness of the impregnating substance (Talc PA), proving that only seed pelletization does not benefit the development of tomato seedlings.

The use of *Natrum muriaticum* dynamizations in tomato seed pelletization favored the performance of GSI and GP, differing statistically from the controls 1 and 2. The treatments pelleted seeds/talc + *Nat-m* 5cH, pelleted seeds/talc + *Nat-m* 7cH, and pelleted seeds/talc + *Nat-m* 9cH had averages higher than the other treatments (Table 3) for both variables.

Table 1. Summary of analysis of variance of germination speed index (GSI), germination percentage (GP), leaf area (LA) and number of leaves (NL) of tomato seedlings from seeds pelleted or not with *Natrum muriaticum* submitted to saline stress.

Variation source	Degree of freedom	Mean squares			
		GSI	GP	LF	NL
Treatments	7	17.46**	1635.26**	4.059**	1.331**
Residues	24	0.28	38.54	1.077	0.032
Variation coefficient (%)		16.93	13.51	26.90	16.56

(**) Significant at 1% probability respectively with f-test.

Table 2. Summary of analysis of variance of root volume (RV), shoot length (SL), shoot dry matter (SDM) and root dry matter (RDM) of tomato seedlings from seeds pelleted or not (control) with *Natrum muriaticum* submitted to saline stress.

Variation source	Degree of freedom	Mean square			
		RV	SL	SDM	RDM
Treatments	7	1.96**	669.68**	0.0044**	0.00099**
Residues	24	0.088	13.79	0.00023	0.000055
Variation coefficient (%)		35.33	3.37	22.45	31.22

(**) Significant at 1% probability respectively with f-test.

Table 3. Mean values of germination speed index (GSI), germination percentage (GP), leaf area (cm², LA) and number of leaves (NL) of tomato seedlings from seeds pelleted or not with *Natrum muriaticum* submitted to saline stress.

TREATMENTS	GSI	GP	LA	NL
Pelleted seeds/ Talc + <i>Nat-m</i> 3cH	1.25 d	25.00 c	3.33 ab	0.62 de
Pelleted seeds/ Talc + <i>Nat-m</i> 5cH	5.75 a	70.00 a	5.22 a	2.12 a
Pelleted seeds/ Talc + <i>Nat-m</i> 7cH	4.81 ab	62.50 ab	4.84 a	1.50 b
Pelleted seeds/ Talc + <i>Nat-m</i> 9cH	4.56 abc	62.50 ab	4.49 ab	1.32 bc
Pelleted seeds/ Talc + <i>Nat-m</i> 11cH	4.437 bc	57.50 ab	3.97 ab	1.20 bc
Pelleted seeds/ Talc + <i>Nat-m</i> 13cH	3.50 c	50.00 b	3.73 ab	0.95 cd
Pelleted seeds/ Talc + distilled water (control 1)	0.54 d	22.50 c	2.10 b	0.47 e
Non-pelleted seeds (control 2)	0.53 d	20.00 c	3.17 ab	0.45 e

Means values followed by at least a same low case letter, in the column, do not differ significantly between themselves by Tukey's test at 1% probability.

Marques-Silva & Bonato [19] reported the action of homeopathic medicinal products of *Rosmarinus officinalis* and *Artemisia absinthium*, in dynamizations 3cH, 6cH, 9cH, 12cH, 24cH, and 30cH, on the germination and growth of common morning glory. The increase in the number of germinated seeds occurred in the 12cH, 24cH, and 30cH dynamizations of *Rosmarinus*.

The use of seed pelletization associated with *Nat-m* 5cH and *Nat-m* 7cH promoted a significant increase by 40.22% and 43.3% in the leaf area (LA) of tomato seedlings grown in saline solution when compared with the control treatment 1 (pelletized seeds/talc + distilled water) (Table 3). According to Oliveira et al. [20], tomato plants irrigated with dam water, with an electrical conductivity of 5.7 dS.m⁻¹, had a 42% reduction in leaf area compared to plants irrigated with

rainwater (0.01 dS.m⁻¹).

Table 4. Means values of root volume (mL, RV), shoot length (mm, SL), shoot dry matter (g, SDM) and root dry matter (g, RDM) of tomato seedlings from seeds pelleted or not (control) with *Natrum muriaticum* submitted to saline stress.

TREATMENTS	RV	SL	SDM	RDM
Pelleted seeds/ Talc + <i>Nat-m</i> 3cH	0.35 cd	16.62 c	0.0521 cde	0.0156 bc
Pelleted seeds/ Talc + <i>Nat-m</i> 5cH	1.87 a	42.22 a	0.1198 a	0.0489 a
Pelleted seeds/ Talc + <i>Nat-m</i> 7cH	1.77 a	39.97 a	0.0972 ab	0.0426 a
Pelleted seeds/ Talc + <i>Nat-m</i> 9cH	1.05 b	37.48 a	0.0846 abc	0.0321 ab
Pelleted seeds/ Talc + <i>Nat-m</i> 11cH	0.97 bc	35.06 ab	0.0830 bc	0.0200 bc
Pelleted seeds/ Talc + <i>Nat-m</i> 13cH	0.47 bcd	28.24 b	0.0617 bcd	0.0184 bc
Pelleted seeds/ Talc + distilled water (control 1)	0.12 d	11.43 c	0.0299 de	0.0075 c
Non-pelleted seeds (control 2)	0.10 d	11.17 c	0.0229 e	0.0061 c

Means values followed by at least a same low case letter, in the column, do not differ significantly between themselves by Tukey's test at 1% probability.

For the NL variable, means were superior in the treatment pelleted seeds/talc + *Nat-m* 5cH, differing statistically from the other treatments. It promoted 22.16% and 21.22%, respectively, in the number of leaves compared to controls 1 and 2 (Table 3). Similar behavior was reported in field conditions in the producing region of Marília, SP. The treatment of passion fruit plants with *Silicea* 30cH promoted a 60% increase in the number of leaves, which later showed a significant correlation with the number of fruits [21].

For the RV, there was a similar behavior to the other variables mentioned above. The treatments pelleted seeds/talc *Nat-m* 5cH and pelleted seeds/talc *Nat-m* 7cH were statistically higher than the other treatments (Table 4). Several researchers have proved homeopathy as input for vegetables under stressful conditions, reporting effective results. The application of *Sulfur*, regardless of potency, was effective in the detoxification of corn plants when exposed to Al³⁺ (150 µM). The treatments *Sulfur* 5cH and *Sulfur* 30cH were effective. *Sulfur* 5cH, 12cH, 30cH, 200cH and 1McH increased by 5.3%, 6.6%, 2.7%, 12.3%, and 6.2%, respectively, the root growth compared to the control [22].

The use of seed pelletization with *Natrum muriaticum* in the 5cH, 7cH, 9cH, 11cH, and 13cH dynamizations also promoted an increased shoot length (SL) of tomato seedlings subjected to saline stress, differing from the 3cH dynamization and the controls 1 and 2 (Table 4).

Pelleted seeds/talc *Nat-m* 5cH, pelleted seeds/talc *Nat-m* 7cH, and pelleted seeds/talc *Nat-m* 9cH were efficient in increasing shoot dry matter (SDM) and root dry matter (RDM) (Table 4). According to Oliveira et al. [20], the variables related to dry biomass represent the most remarkable effects of salinity on tomato plants.

Mechanisms and interactions were also observed in dynamized systems, such as detoxification mechanisms in seeds and plants intoxicated with heavy metals, as reported for plants of *Pisum sativum* (pea) intoxicated by copper sulphate and treated with *Cuprum sulphuricum* 15cH, a

phenomenon also noted for *Cuprum sulphuricum* 14cH in *Sinapis alba* var. *Dialba* (mustard), and *Cuprum sulphuricum* 5cH in *Triticum aestivum* (wheat) [23-26].

Studies with arsenic in seeds of *Triticum aestivum* (wheat) previously subjected to lethal doses of arsenic (physiological stress) and later treated with dynamizations of arsenic trioxide (As₂O₃) 40x, 42x, 45x, non-dynamized As₂O₃, distilled water, and dynamized distilled water 40x, 42x, 45x demonstrated the most promising effects on the dynamization of As₂O₃ 45x and on stress. However, the decimal dynamizations of water were more significant when compared to the control (distilled water), and the high dilutions of non-dynamized As₂O₃ were not effective [27-29].

The development of tomato seedlings from seeds pelleted with *Natrum muriaticum* 5cH and 7cH was better for all evaluated variables, differing statistically from the controls, confirming the positive response of homeopathy in the development of seedlings under disequilibrium conditions.

Conclusion

Natrum muriaticum 5cH and 7cH cause an increase in the production of healthy tomato seedlings submitted to saline stress.

The process of seed coating using homeopathic preparations is a promising technology for applying homeopathy to vegetables.

References

- [1] FAO. Water in agriculture: opportunity untapped. Rome (Italy): Food and Agriculture Organization of the United Nations; 2006.
- [2] Holanda JS, Amorim JRA, Ferreira Neto M, Holanda AC. Qualidade da água para irrigação. In: Gheyi HR, Dias NS, Lacerda CF, editors. Manejo da salinidade na Agricultura: estudos básicos e aplicados. Fortaleza (Brazil): INCTSal; 2010. p. 43-61.
- [3] Taiz L, Zeiger E. Fisiologia Vegetal. 3rd ed. Porto Alegre (Brazil): Artmed; 2004.
- [4] Cavalcante LF. Sais e seus problemas nos solos irrigados [dissertation]. Areia (Brazil): Center of Agricultural Sciences, Federal University of Paraíba; 2000. 72p.
- [5] Doorenbos J, Kassam AH. Efeito da água no rendimento das culturas. Gheyi HR, Sousa AA, Damasceno FAV, Medeiros JF, translators. Campina Grande (Brazil): UFPB; 1994. (FAO. Estudos de Irrigação e drenagem, 33).
- [6] Blanco FF. Tolerância do tomateiro à salinidade sob fertirrigação e calibração de medidores de íons específicos para determinação de nutrientes na solução do solo e na planta. [thesis (doctorate)]. Piracicaba (Brazil): College of Agriculture "Luiz of Queiroz", University of São Paulo; 2004. 115p.
- [7] Araújo Filho JB, Gheyi HR, Azevedo NC. Tolerância da bananeira à salinidade em fase inicial de desenvolvimento. Pesquisa Agropecuária Brasileira. 1995; 30(7): 989-997. [portuguese].

- [8] Silva HA, Parizotto AV, Moreira FC, Marques RM, Reis B, Bonato CM. The effect of high dilutions of *Pulsatilla nigricans* on the vigour of soybean seeds subjected to accelerated aging. *Acta Scientiarum. Agronomy*. 2012; 34(2): 201-206.
- [9] Bonfim FPG, Martins ER, Dores RGR, Barbosa CKR, Casali VWD, Honório ICG. Use of homeopathic *Arnica montana* for the issuance of roots of *Rosmarinus officinalis* L. and *Lippia alba* (Mill) N.E.Br. *Int J High Dilution Res* [online]. 2008 [cited 2019 Apr 15]; 7(23); 113-117. Available from:
<http://journal.giri-society.org/index.php/ijhdr/article/view/276/348>
- [10] Almeida AA, Galvão JCC, Casali VWD, Lima ER, Miranda GV. Tratamentos homeopáticos e densidade populacional de *Spodoptera frugiperda* (J.E.Smith,1797) (Lepdoptera: Noctuidae) em plantas de milho no campo. *Revista Brasileira de Milho e Sorgo*. 2003; 2(2): 32-39. [portuguese].
- [11] Toledo MV, Stangarlin JR, Bonato CM. Controle da pinta preta em tomateiro com preparados homeopáticos de própolis. *Revista Brasileira de Agroecologia*. 2009; 4(2): 471-474. [portuguese].
- [12] Andrade FMC, Casali VWD, de Vita B, Cecon PR, Barbosa LCA. Efeito de homeopatas no crescimento e na produção de cumarina em chambá (*Justicia pectoralis* Jacq.). *Revista Brasileira de Plantas Medicinai*s. 2001; 4(1): 19-28. [portuguese].
- [13] Almeida MAZ. Resposta do manjeriçao (*Ocimum basilicum* L.) à aplicação de preparações homeopáticas [dissertation]. Viçosa (Brazil): Department of Phytotechnics, Federal University of Viçosa; 2002. 112p.
- [14] Arruda VM. Aplicações de soluções homeopáticas em *Achilleia millefolium* L. (Asteraceae): abordagem morfofisiológica [dissertation]. Viçosa (Brazil): Department of Phytotechnics, Federal University of Viçosa; 2005. 107p.
- [15] Casali VWD, Andrade FMC, Duarte ESM. Acologia de altas diluições: resultados científicos e experiências sobre uso de preparados homeopáticos em sistemas vivos. Viçosa (Brazil): Department of Phytotechnics, Federal University of Viçosa; 2009.
- [16] Brazilian Homeopathic Pharmacopoeia. 2nd ed. São Paulo: Atheneu, 1997.
- [17] Fernandes AA, Martínez HEP, Oliveira LR. Produtividade, qualidade dos frutos e estado nutricional de plantas de pepino, cultivadas em hidroponia, em função das fontes de nutrientes. *Horticultura Brasileira*. 2002; 20(4): 571-575. [portuguese].
- [18] Maguire JD. Speed of germination-aid in selection and evaluation for seedling emergence and vigor. *Crop Science*. 1962; 2(1):176-177.

- [19] Marques-Silva GG, Bonato CM. Avaliação da aplicação das soluções homeopáticas de *Rosmarinus officinalis* L. e *Artemisia absinthium* L. na germinação e no desenvolvimento de corda-de-viola (*Ipomea* sp.). Cultura Homeopática. 2006; 16(5): 50-50. [portuguese].
- [20] Oliveira BC, Cardoso MAA, Oliveira JC, Oliveira FA, Cavalcanti LF. Características produtivas do tomateiro submetido a diferentes níveis de sais, na água de irrigação. Rev. Bras. Eng. Agríc. Ambient. 2007; 11(1): 11-16. [portuguese].
- [21] Rolim PRR, Brignani Neto F, Souza SA, Mizote FA, Narita N, Jesus CR, Shinohara D, Oliveira DA. Manejo da cultura de maracujá sem o uso de agroquímicos convencionais. Proceedings of the IV Reunião Técnica de Pesquisa em Maracujazeiro; 2002; Viçosa (Brazil): Federal University of Viçosa; 2002. p.113. [portuguese].
- [22] Rocha M, Moretti MR, Bonato CM. Efeito de diferentes dinamizações homeopáticas de *Sulphur* no comprimento da raiz principal de plântulas de milho (*Zea mays*) sob estresse por alumínio. Proceedings of the XI Encontro Anual de Iniciação Científica; 2002 Oct 01-04; Maringá (Brazil): State University of Maringá; 2002. [portuguese].
- [23] Nieten G, Boiron J, Marin A. Ação de doses infinitesimais de sulfato de cobre sobre plantas previamente intoxicadas por essa substância; ação da 15ª centesimal hahnemanniana. Pesquisa experimental moderna em Homeopatia. Rio de Janeiro (Brazil): Editorial Homeopática Brasileira; 1969.
- [24] Auquièrre JP, Moens P. Recherches de l'action de dilutions homéopathiques sur les végétaux. I. Action du CuSO_4 5CH sur blé intoxiqué au CuSO_4 1%. Journal de pharmacie de Belgique. 1981; 36(1): 303-320. [french].
- [25] Auquièrre, JP, Moens P, Martin PL. Recherche de l'action de dilutions homéopathiques sur les végétaux. II. Action du CuSO_4 14DH sur la Moutarde blanche (*Sinapis alba* L.; var *dialba*) intoxiquée au CuSO_4 0.1 et 0.2%. Journal de pharmacie de Belgique. 1982; 37(1): 117-134. [french].
- [26] Scolfield AM. Homoeopathy and its potential role in agriculture – a critical review. Biological Agriculture and Horticulture. 1984; 2(1):1-50.
- [27] Brizzi M, Betti L, Nani D, Peruzzi M. An overall analysis of a series of experiments based on high dilutions in an *Arsenicum album*/wheat modelo. Omeomed. 1997; 97(1): 12-13.
- [28] Brizzi M, Nani D, Peruzzi M, Betti L. Statistical analysis of the effect of high dilutions of arsenic in a large dataset from a wheat germination model. British Homeopathic Journal. 2000; 89(2): 63-67.

[29] Brizzi M, Lazzarato L, Nani D, Borghini F, Peruzzi M, Betti L. A biostatistical insight into the As₂O₃ high dilution effects on the rate and variability of wheat seedling growth. *Forschende Komplementärmedizin und Klassische Naturheilkunde*. 2005; 12(5): 277-283.

Received: April 18, 2019. Accepted: June 6, 2019

© International Journal of High Dilution Research.
Not for commercial purposes.

